

Attachment: The designated diagnostic tests and other quarantine measures required for designated mammals

(In case of any discrepancy between the English version and the Chinese text, the Chinese text shall govern.)

Document No.: 1121481435 (20230323)

Scientific name	Chinese name	The animals must undergo the designated diagnostic tests during the pre-export quarantine period with negative results and meet the required quarantine measures.
<i>Rodentia</i>	齧齒動物	<ol style="list-style-type: none"> 1. For leptospirosis: microscopic agglutination test; 2. For toxoplasmosis: competitive enzyme-linked immunosorbent assay (c-ELISA) or the polymerase chain reaction (PCR) test.
<i>Dasypodidae</i>	犛狽科動物	<ol style="list-style-type: none"> 1. For leptospirosis: microscopic agglutination test; 2. For toxoplasmosis: c-ELISA or PCR test.
<i>Choloepus didactylus</i>	二趾樹懶	For leptospirosis: microscopic agglutination test.
<i>Tamandua tetradactyla</i>	食蟻獸	For toxoplasmosis: c-ELISA or PCR test.
<i>Camelidae</i>	駱駝科動物	<ol style="list-style-type: none"> 1. Animals must be raised in establishments which comply with the following requirements: <ol style="list-style-type: none"> 1.1. No cases of the following diseases have been confirmed in the establishments of origin in the previous two years: tuberculosis (<i>Mycobacterium bovis</i>, <i>M. caprae</i>, and <i>M. tuberculosis</i>), paratuberculosis (Johne's disease), <i>Echinococcus granulosus</i> infections,

		<p>anaplasmosis, babesiosis, and theileriosis.</p> <p>1.2. No cases of the following diseases have been confirmed in the establishments of origin in the previous year: rabies, anthrax, bluetongue, alpaca fever (caused by <i>Streptococcus pyogenes</i>), peste des petits ruminants (PPR), Rift Valley fever, and Q fever.</p> <p>1.3. No cases of the following diseases have been confirmed in the establishments of origin in the previous six months: camel pox, contagious pustular dermatitis (ORF), West Nile fever, epizootic hemorrhagic disease (caused by the <i>Reoviridae</i> virus), bovine viral diarrhea, leptospirosis, equine herpesvirus-1 infections, brucellosis (caused by <i>Brucella abortus</i> and <i>B. melitensis</i>), equine encephalomyelitis (Eastern and Western), and Surra (caused by <i>Trypanosoma evansi</i>).</p> <p>2. The designated diagnostic tests:</p> <p>2.1. For tuberculosis: Intradermal tuberculin test;</p> <p>2.2. For brucellosis (<i>Brucella abortus</i>, <i>B. melitensis</i> and <i>B. suis</i>): c-ELISA, complement fixation test (CF), serum tube agglutination test, or Rose-Bengal test;</p> <p>2.3. For bluetongue: c-ELISA;</p> <p>2.4. For paratuberculosis (Johne's disease): ELISA;</p> <p>2.5. For bovine viral diarrhea: antigen detection ELISA;</p>
--	--	--

		<p>2.6. For Q fever: CF or ELISA;</p> <p>2.7. For epizootic hemorrhagic disease: c-ELISA;</p> <p>2.8. For bovine anaplasmosis: card agglutination test or c-ELISA;</p> <p>2.9. For middle east respiratory syndrome coronavirus (MERS-CoV) infection, only for <i>Camelus dromedaries</i>: PCR.</p> <p>3. Alpacas and llamas must be sheared (whole body, including the head and lower legs) within 60 days prior to export.</p>
--	--	--