

## Attachment: The designated diagnostic tests and other quarantine measures required for designated mammals

(In case of any discrepancy between the English version and the Chinese text, the Chinese text shall govern.)

Document No.: 1121481695 (20230419)

| Scientific name    | Chinese name | The animals must undergo the designated diagnostic tests during the pre-export quarantine period with negative results and meet the required quarantine measures.   |
|--------------------|--------------|---|
| <i>Rodentia</i>    | 齧齒動物         | <ol style="list-style-type: none"> <li>1. For leptospirosis: microscopic agglutination test. If the required tests for leptospirosis are not conducted, then all animals must be administered with penicillin or other medication deemed as of equivalent efficacy in treating or preventing leptospirosis by the animal quarantine authority of exporting country;</li> <li>2. For toxoplasmosis: enzyme-linked immunosorbent assay (ELISA) or the polymerase chain reaction (PCR) test. If the required tests for toxoplasmosis are not conducted, the animals must be administered with pyrimethamine, or other medication deemed as of equivalent efficacy in treating or preventing toxoplasmosis by the animal quarantine authority of the exporting country</li> </ol> |
| <i>Dasypodidae</i> | 犴狢科動物        | <ol style="list-style-type: none"> <li>1. For leptospirosis: microscopic agglutination test;</li> <li>2. For toxoplasmosis: ELISA or PCR test.</li> </ol>   |
| <i>Choloepus</i>   | 二趾樹懶         | For leptospirosis: microscopic agglutination test.  |

|  |       |   |
|--|-------|---|
| <i>didactylus</i>                      |       |   |
| <i>Tamandua</i><br><i>tetradactyla</i> | 食蟻獸   | For toxoplasmosis: ELISA or PCR test.   |
| <i>Camelidae</i>                       | 駱駝科動物 | <ol style="list-style-type: none"> <li>1. Animals must originate from a state or province where vesicular stomatitis, caprine and ovine brucellosis, and contagious caprine pleuropneumonia have not occurred in the previous year.</li> <li>2. Animals must be raised in establishments which comply with the following requirements: <ol style="list-style-type: none"> <li>2.1. No cases of the following diseases have been confirmed in the establishments of origin in the previous two years: tuberculosis (<i>Mycobacterium bovis</i>, <i>M. caprae</i>, and <i>M. tuberculosis</i>), <i>Echinococcus granulosus</i> infections, anaplasmosis, babesiosis, and theileriosis.</li> <li>2.2. No cases of the following diseases have been confirmed in the establishments of origin in the previous year: rabies, anthrax, bluetongue, paratuberculosis (Johne's disease), alpaca fever (caused by <i>Streptococcus pyogenes</i>), peste des petits ruminants (PPR), Rift Valley fever, and Q fever.</li> <li>2.3. No cases of the following diseases have been confirmed in the establishments of origin in the previous six months: scrapie, camel pox, contagious pustular dermatitis (ORF), West</li> </ol> </li> </ol> |

|  |  |  |
|--|--|--|
|  |  | <p>Nile fever, epizootic hemorrhagic disease (caused by the <i>Reoviridae</i> virus), bovine viral diarrhea, leptospirosis, equine herpesvirus-I infections, equine encephalomyelitis (Eastern and Western), and Surra (caused by <i>Trypanosoma evansi</i>).</p> <p>3. The designated diagnostic tests:</p> <p>3.1. For tuberculosis: Intradermal tuberculin test;</p> <p>3.2. For caprine and ovine brucellosis (<i>Brucella abortus</i> and <i>B. melitensis</i>): ELISA, complement fixation test (CF), serum tube agglutination test, Rose-Bengal test or Buffered <i>Brucella</i> antigen test;</p> <p>3.3. For bluetongue: c-ELISA;</p> <p>3.4. For paratuberculosis (Johne's disease): CF or ELISA;</p> <p>3.5. For bovine viral diarrhea: antigen detection ELISA;</p> <p>3.6. For Q fever: CF or ELISA;</p> <p>3.7. For epizootic hemorrhagic disease: c-ELISA;</p> <p>3.8. For bovine anaplasmosis: CF, card agglutination test or ELISA;</p> <p>3.9. For middle east respiratory syndrome coronavirus (MERS-CoV) infection, only for <i>Camelus dromedaries</i>: PCR.</p> <p>4. Animals must be injected with long-acting oxytetracycline at a dose rate of 20 mg/kg during the pre-export quarantine period</p> <p>5. Alpacas and llamas must be sheared (whole</p> |
|--|--|--|

|  |  |  |
|--|--|--|
|  |  | body, including the head and lower legs) within 60 days prior to export. |
|--|--|--|